

Risk of radiation exposure of the lens of the eye for members of the public Bibliography study

M.-A. Chevallier, A. Rannou, I. Clairand, C. Villagrasa

Institut de Radioprotection et de Sûreté Nucléaire, IRSN



Laboratory of Ionizing Radiation Dosimetry

Bât 02 - BP 17 - 31, avenue de la Division Leclerc, 92260 Fontenay-aux-Roses - France

Email : marie-anne.chevallier@irsn.fr

Objectives of this literature review

New dose limit recommended by ICRP for the lens of the eye for occupational exposure : \setminus to 20 mSv per year [1] No explicit recommendation regarding the dose limit for the public

Are there potential situations where public exposure can lead to equivalent doses to the lens of the eye exceeding 1mSv per year?

Methodology

Three steps:

1/ Exhaustive list of consumer goods containing radioactive substances [2] [3]

17 Exhibitive for Consumer goods contaming radiocates assumed [5] [6]
27 Focus on goods that may lead to significant eye lens exposure
37 IRSN calculations of dose to the lens of the eye using when necessary MCNPX code [5]

Scenarii chosen in order to reflect daily life situations

Calculations made with the spare quantitative information from the literature review

Situations for workers and patients are not considered in the scope of this study

Weapon sights

Tritium used to illuminate the aiming triangle in the night

Not widely spread, concerns only specialists Radionuclides: tritium (up to 10 GBq)

Photographic films

Old photographic prints, developed using uranium nitrate

Exotic situation Radionuclides: uranium

No risk of significant radiation exposure to the eye lens

TV receivers and PC monitors

Television receivers or PC monitors with cathode-ray tubes

Essentially television manufactured before 1970 X-rays (up to 5.3 µSv.h·¹ averaged over 10 cm² at any readily accessible point 5 cm from the surface)



B

Timepieces

Radioluminous paint added to timepieces

Not widely spread, concerns only amateur repairing timepieces

Radionuclides: radium-226 (170 kBq maximum), tritium or promethium-147

~0.25 mSy per year Not significant

Limited or even insignificant radiation exposure to the eye lens

Naturally radioactive minerals

Concerns only amateur collecting radioactive minerals

Radionuclides: uranium, thorium, 40K, radium-226

For photon radiations only: ~ 0.85 mSv per year -2.3 mSv per year (if 1 ppm of ²²⁶Ra) 0.01 μSv per year Complete calculation in progress

Significant exposure for some minerals, but very rare situations

Dental products

Dental ceramics with added uranium

Before the 1980's, natural or depleted uranium was added in dental porcelains for aesthetic purpose (fluorescent properties)

Only few people may be concerned

Radionuclides: uranium (≤ 10 kBg.g-1), 40K

Dose to the eye lens induced by γ radiations only ≤ 0.1 mSv

New dental ceramics

Nowadays, only natural presence of potassium-40, rubidium-87, and uranium and thorium (as impurities)

Radionuclides: uranium, thorium, 40K, 47Ru

Amounts of several orders of magnitude below former dental ceramics

* Beta emitter at low energy

1400

Activities similar to the former dental ceramics

No risk of significant radiation exposure to the eye lens

Gemstones

Two scenarii of exposure: gemstones in a piercing (0.02 carat, 3 cm from the eye) or in earrings (2x10 carat, 7 cm from the eye)

Naturally radioactive gemstones

Uranium, thorium and potassium-40 may be naturally present in some gemstones as zircon (widely used) or eukanite, thorianite,...

- Zircon: ≤ 7 µSv per year → Dose to the eye lens is not significant
- Other gemstones: up to 31 mSv per year **Exotic situations** (143 kBg of natural thorium in thorianite)

Irradiated gemstones

Some gemstones are irradiated to modify their colour Most commonly irradiated gemstones : topaz also: tourmaline, zircon, beryl, quartz, diamonds...

Many activated elements due to the irradiation ⁴⁶Sc, ⁵⁴Mn, ¹³⁴Cs, ... and thorium (as impurities)



- Concentrations at higher activity > Very rare situations

Lenses

Thorium occurs naturally in rare earths and zirconium oxides used to process the glass, or thorium is intentionally added to improve optical properties. It can also be used in surface coating.

Lenses in eveglasses

500 kBg of natural thorium Eyeglasses worn 16 hours per day, all the year

Dose rate: 0.3 mSv.h-1

1 mSv/year is reached for activity ~ 0.3 kBq

Significant dose rate but the activity of thorium considered could not be confirmed

Lenses in cameras

500 kBg of natural thorium Viewfinder of a camera, used 1 h per week, all the year

Dose rate: 2.8 µSv.h-1

1 mSv/year is reached for activity ~7 kBq

Very rare situations



Conclusion

No current situation was identified where member of the public might receive significant radiation dose to the lens of the eye.

The very few existing situations where the dose to the lens of the eye could reach 1 mSv/year are exotic ones:

- One significant exposure comes from ophthalmic lenses with 500 kBq of thorium. But this activity mentioned in the literature could not be confirmed and no information about the use of those items nor their quantity on the marketplace were available.
- The second situation comes from radioactive minerals that could be owned by mineral collectors, and especially radiobarites. The highest concentration of radium-226 found in the bibliography could not be confirmed, and the only other values of concentrations found in the literature review are several orders of magnitude lower. In that case, there is no significant exposure to the lens of the eye any more.

REFERENCES

ational Commission on Radiological Protection, Statement on Tissue Reactions, approved by the Commission on April 21, 2011
w, J. Dunderdale and R. A. Paynter, RADIATION PROTECTION 146, A Review of Consumer Products Containing Radioactive Substances in the European Union, NRPB Occupational Services Department, 2007 and Council on Radiation Protection, Radiation Exposure of the U.S. Population from Consumer Products and Council on Reciliance Sources, NCRP Report No. 95, 1987
Chevaliter, A. Rannou, I. Clairand, C. Villagrasa, Risk of radiation exposure of the lens of the eye for members of the public, bibliography study, IRSN report, PRP-HOM/SDE n° 2012-00061
endricks et al. MCNPX 2.6.0 Extensions, LA-UR-08-02-216, April 2005